


# Chemistry Notes for class 12 Chapter 12

## Aldehydes, Ketones and Carboxylic Acids

In aldehydes, the carbonyl group ( $\text{C}=\text{O}$ ) is bonded to carbon and hydrogen, while in the ketones, it is bonded to two carbon atoms

### Nature of Carbonyl Group

The carbon and oxygen of the carbonyl group are  $\text{Sp}^2$  hybridised and the carbonyl double bond contains one  $\sigma$ -bond and one  $\pi$ -bond.



The electronegativity of oxygen is much higher than that of the carbon, so there electron cloud is shifted towards the oxygen. Therefore, C-O bond is polar in nature.

### Nomenclature

(i) **Nomenclature of aldehydes** In IUPAC system, the suffix “e” of alkane is replaced by the suffix “al”. e.g.,


Compound	Common name	IUPAC name
HCHO	Formaldehyde	Methanal
$\text{CH}_3\text{CHO}$	Acetaldehyde	Ethanal

(ii) **Nomenclature of ketones** In IUPAC system, the suffix “e” of alkane is replaced by “one”. e.g.,


Compound	Common name	IUPAC name
$\text{H}_3\text{C} \cdot \text{COCH}_3$	Dimethyl ketone (acetone)	Propanone
$\text{H}_3\text{C} \cdot \text{COC}_2\text{H}_5$	Ethyl methyl ketone	Butanone

### Preparation of Aldehydes and Ketones


(i) **By oxidation of alcohols** Aldehydes and ketones are generally prepared by oxidation of primary and secondary alcohols, respectively.




(ii) **By dehydrogenation of alcohols** In this method, alcohol vapours are passed over heavy metal catalysts (Ag or Cu). Primary and secondary alcohols give aldehydes and ketones.




(iii) **By ozonolysis of alkenes**






(iv) **By hydration of alkynes** Acetylene on hydration gives acetaldehyde and other alkynes on hydration give ketones.



**(v) By heating Ca salt of acid**

To obtain aldehyde, calcium formate and any other Ca salt of acid are heated.

**(vi) By decarboxylation and dehydration of aromatic acids****Preparation of Aldehydes**


**(i) Rosenmund reduction**

Formaldehyde cannot be prepared by this method as  $\text{HCOCl}$  is highly unstable.



**(ii) From nitriles**

This reaction is called **Stephen reaction**.


Alternatively, nitriles are selectively reduced by diisobutylaluminium hydride, [DiBAL-H] to imines which on hydrolysis give aldehydes.




Similarly, esters can also be reduced to aldehydes with DiBAL-H.

**(iii) Etard reaction**





(iv) **Side chain halogenation followed by hydrolysis of toluene**



(v) **Gattermann-Koch synthesis**



**Preparation of Ketones**

**(i) From acyl chlorides****(ii) From nitriles****(iii) Friedel-Crafts acylation****(iv) Oppenauer oxidation****Physical Properties of Aldehydes and Ketones**

1. Methanal (HCHO) is a gas at room temperature, and its 40% aqueous solution is known as formalin. It is a reducing agent in silvering of mirrors and decolourising vat dyes.
2. Ethanal (CH<sub>3</sub>CHO) is a volatile liquid. Other aldehydes and ketones are liquid or solid at room temperature.
3. The boiling point of aldehydes and ketones are higher than hydrocarbons and ethers of comparable molecular mass due to high magnitude of dipole-dipole interactions.
4. Aldehydes and ketones have lower boiling point than those of alcohols of similar molecular


masses due to absence of intermolecular hydrogen bonding.

5. The lower members of aldehydes and ketones are miscible with water due to the formation of hydrogen bond with water. However, the solubility decreases with increase in length of alkyl chain.


6. Acetophenone is a hypnotic (sleep producing drug) so used as a medicine under the name hypnone.

## Chemical Reactions of Aldehydes and Ketones

### (i) Nucleophilic addition reactions



Reactivity order is




It is due to +I effect of alkyl groups which decreases the positive charge on carbonyl carbon and steric hinderance (The bulky alkyl group hinder the approach of nucleophile).

Addition of hydrogen cyanide




Addition of sodium hydrogen sulphite




This reaction is used for the separation and purification of aldehydes and ketones. This is because the addition compound formed, is water soluble and can be converted back to the original carbonyl compound by treating with dilute mineral acid or alkali.

Addition of lower alcohols





(ii) **Addition of ammonia and its derivatives** Reaction with ammonia






Urotropine on controlled nitration gives the well known explosive RDX (Research and Development explosive).






### Reaction with ammonia derivatives




where, Z = alkyl, aryl, —OH, —NH<sub>2</sub>, —C<sub>6</sub>H<sub>5</sub>NH, —NHCONH<sub>2</sub> etc.

### Some N-substituted Derivatives of Aldehydes and Ketones


Z	Reagent name	Carbonyl derivative	Product name
—R	Amine	$>C=N-R$	Substituted imine (Schiff's base)
—OH	Hydroxyl amine	$>C=N-OH$	oxime
—NH <sub>2</sub>	Hydrazine	$>C=N-NH_2$	Hydrazone
—NH— 	Phenyl-hydrazine	$>C=N-NH-$	Phenylhydrazone
—NH— 	2,4-dinitro-phenyl hydrazine	$>C=N-NH-$	2,4-dinitro-phenyl hydrazone
—NH— 	Semi-carbazide	$>C=N-NH-$	Semi-carbazone

(iii) **Reduction** Aldehydes and ketones are reduced to primary and secondary alcohols respectively by sodium borohydride (NaBH<sub>4</sub>) or lithium aluminium hydride [LiAlH<sub>4</sub>].


#### Clemmensen reduction




#### Wolff-Kishner reduction



(iv) **Oxidation** Aldehydes get easily oxidised to carboxylic acids by HNO<sub>3</sub>, KMnO<sub>4</sub>, K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>, etc., or even by mild oxidising agent.




**Ketones** are generally oxidised under vigorous conditions, i.e., strong oxidising agents and at elevated temperature.



During oxidation of unsymmetrical ketones the point of cleavage is such that keto group stays preferentially with the smaller alkyl group (popoff's rule).

(a) **Tollen's test** Aldehydes give bright silver mirror with Tollen's reagent (ammoniacal silver nitrate).



(b) **Fehling's test** Fehling solution gives a reddish brown precipitate with aldehydes (except benzaldehyde)




[Fehling solution is a mixture of Fehling solution A and Fehling solution B in 1: 1 ratio. Fehling solution A is aqueous copper sulphate and Fehling solution B is alkaline sodium potassium tartrate which is also called, Rochelle salt.]


(c) **Benedict solution** With it, aldehydes (except benzaldehyde) also give red ppt. of Cu<sub>2</sub>O.

(d) **Schiff's reagent** It is an aqueous solution of magenta or pink coloured rosaniline hydrochloride which has been decolourised by passing SO<sub>2</sub>. Aldehydes give pink colour with this reagent but ketones do not.

**Haloform reaction** Aldehydes and ketones having at least one methyl group [3-α hydrogen] linked to the carbonyl carbon atom (methyl ketones) are oxidised by sodium hypohalite to sodium salts of corresponding carboxylic acids having one carbon atom less than that of carbonyl compound. The methyl group is converted to haloform.




This oxidation does not affect a carbon-carbon double bond, if present in the molecule.



Iodoform reaction with sodium hypoiodite is also used for the detection of  $\text{CH}_3-$  group or  $\text{CH}_3\text{CH}(\text{OH})-$  group by producing yellow solid  $\text{CHI}_3$ .




#### (v) Aldol condensation



Its further condensation gives phorone,


This reaction is exhibited by those aldehydes and ketones which have at least one  $\alpha$ -hydrogen.

(vi) **Cross aldol condensation** Base catalysed crossed aldol condensation between an aromatic aldehyde and an aliphatic aldehyde or ketone is called **Claisen-Schmidt condensation** or **Claisen reaction**.




The above reaction is called Benzoin condensation, not the cross aldol condensation.


(vii) **Cannizzaro reaction** Aldehydes which do not have any  $\alpha$  – hydrogen atom, undergo self oxidation and reduction (disproportionation) reaction on treatment with concentrated alkali.




(viii) **Electrophilic substitution reaction** Aromatic aldehydes and ketones undergo electrophilic substitution. Carbonyl group shows + R effect, therefore acts as a deactivating and meta directing group.






(ix) **Baeyer- Villiger oxidation** With Caro's acid ( $\text{H}_2\text{SO}_5$ ) or per benzoic acid ( $\text{C}_6\text{H}_5\text{CO}_3\text{H}$ ) or peracetic acid ( $\text{CH}_3\text{CO}_3\text{H}$ ) aliphatic ketones give ester.






(x) **Tischenko's reaction** It is a modified form of Cannizzaro reaction.



(xi) **Knoevenagel reaction** It involves condensation between active methylene group and carbonyl groups in the presence of base.

**(xii) Schmidt reaction****(xiii) Reformatsky reaction**

**(xiv) Perkin's reaction****(xv) Wittig reaction****(xvi) Polymerisation****Carboxylic Acids**

These are the compounds which have  $\begin{array}{c} \text{O} \\ \parallel \\ \text{—C—OH} \end{array}$  group [carboxyl group]. The word carboxyl is a combination of two words carbonyl ( $\text{>C=O}$ ) and hydroxyl ( $\text{—OH}$ ).

**Classification**

Depending upon the number of  $\text{—COOH}$  groups, they are classified as

- (i) monocarboxylic acids; containing one  $\text{—COOH}$  group
- (ii) dicarboxylic acids: containing two  $\text{—COOH}$  groups.


**Sources of carboxylic acids**




Formula	Common name	Source
HCOOH	Formic acid	Red ant (formica)
CH <sub>3</sub> COOH	Acetic acid	Vineger (acetum)
C <sub>3</sub> H <sub>7</sub> COOH	Butyric acid	Butter (butyrum)

## Nomenclature


Their IUPAC names have been derived from the corresponding alkanes by replacing the letter 'li' of the alkane with 'oic' and adding suffix 'acid' at the end, Thus, monocarboxylic acids are called alkanolic acids.



**Benzene carboxylic acid  
(Benzoic acid)**





**2-phenylethanoic acid**




**Benzene-1, 2-dicarboxylic acid  
(phthalic acid)**

## Methods of Preparation of Monocarboxylic Acids


### (i) From primary alcohols and aldehydes



(ii) **From alkyl benzenes** Alkyl benzene when treated with strong oxidising agent like H<sub>2</sub>CrO<sub>4</sub> (chromic acid), acidic or alkaline KMnO<sub>4</sub> gives benzoic acid.




(iii) **From acid derivatives** All acid derivatives like amides ( $\text{RCONH}_2$ ), acid halides ( $\text{RCOCl}$ ), esters ( $\text{RCOOR}'$ ), acid anhydrides ( $\text{RCO-O-COR}$ ) on hydrolysis give carboxylic acids. All acid derivatives break from  $\text{RCO}^+$ .




$Z = -\text{NH}_2, -\text{X} (\text{X} = \text{Cl, Br, I}), \text{OR}', \text{RCOO}-$  etc.

Ease of hydrolysis :  $\text{RCOCl} > (\text{RCO})_2\text{O} > \text{RCOOR}' > \text{RCONH}_2$


(iv) **From nitriles and amides** Nitriles are hydrolysed to amides and then to acids in the presence of  $\text{H}^+$  or  $\text{OH}^-$  as catalyst. Mild reaction conditions are used to stop the reaction at the amide stage.




(v) **From Grignard reagents** Grignard reagents react with carbon dioxide (dry ice) to form salts of carboxylic acids which in turn give corresponding carboxylic acids after acidification with mineral acid



(vi) **By heating geminal dicarboxylic acids**




(vii) **From alkynes**



### Physical Properties of Carboxylic Acids


1. Aliphatic carboxylic acids up to nine carbon atoms are colourless liquids at room temperature with unpleasant odours. The higher acids are wax like solids.
2. The lower carboxylic acids are freely miscible with water due to the presence of intermolecular hydrogen bonding with H<sub>2</sub>O molecules. However, the solubility in water decreases gradually due to increase in the size of alkyl group.
3. Monocarboxylic acids have higher boiling points as compared to the alcohols of comparable molecular masses due to the presence of stronger intermolecular hydrogen bonding as shown below.




4. Melting points of aliphatic monocarboxylic acids shows alternation or oscillation effect, i.e., the m.p. of an acid with even number of carbon atoms is higher than the next lower and next higher homologue containing odd number of carbon atoms. This is because, in case of acids with even number of carbon atoms, the terminal -CH<sub>3</sub> and -COOH groups lie on the opposite sides of the zig-zag chain. As a result, they get closely packed in the crystal lattice.
5. Glacial acetic acid is completely pure acetic acid and represents the solid state of acetic acid. Below 16.6°C temperature pure acetic acid is converted into ice like solid hence it is called glacial acetic acid.

### Chemical Properties of Carboxylic Acids





This is because -I effect decreases in the order : F > Cl > Br > I.




This is because -I effect decreases with distance.


Peracetic acid ( $\text{CH}_3\text{COOOH}$ ) is a weaker acid than acetic acid as acetate ion is stabilised by resonance.

Acidic strength of aromatic acids The parent member of the family benzoic acid which is a weaker acid ( $K_a = 6.3 \times 10^{-5}$ ) than acid ( $K_a = 17.7 \times 10^{-5}$ ) but stronger than acetic acid.

Some order of acidity are




(b) Similarly,  $K_a$  values of methyl substituted (toluic acids) at 298 K are as follows:




From the  $K_a$  values, it is evident that with the exception of o-isomer, both p and m-toluic acids are weaker acids than benzoic acid whereas the three isomeric nitro benzoic acids are stronger acids than benzoic acid.

## (ii) Reactions involving cleavage of C-O-H bond


### (a) Formation of anhydride





### (b) Esterification




## Mechanism




### (c) Reactions with $\text{PCl}_5$ , $\text{PCl}_3$ and $\text{SOCl}_2$



### (iii) Chemical reactions involving – COOH group


**(a) Reduction****(b) Decarboxylation**

(iv) **Substitution reactions in the hydrocarbon part**  $\alpha$ -hydrogen atoms in carboxylic acids are acidic in nature and can be easily replaced by halogen atoms in HVZ reaction




The reaction is known as Hell-Volhard-Zelinsky reaction.

(v) **Arndt-Eistert reaction** It is method of converting lower carboxylic acids to their higher homologues




(vi) **Reducing property** Among carboxylic acids, formic acid is the only acid that acts as reducing agent. It reduces, acidified  $\text{KMnO}_4$  to  $\text{MnSO}_4$ ,  $\text{HgCl}_2$  to  $\text{Hg}$ , Tollen's reagent to silver mirror and Fehling's solution to red ppt. and itself gets oxidised to  $\text{CO}_2$  and  $\text{H}_2\text{O}$ .



(vii) **Electrophilic substitution reactions of aromatic acids**  $-\text{COOH}$  group shows  $-\text{R}$  effect, therefore, acts as a deactivating and meta-directing group. Carboxylic acids do not undergo Friedel-Craft's reaction because the carboxylic group is deactivating and the catalyst  $\text{AlCl}_3$  (anhy.) gets bonded to the carboxyl group.







## Uses

1. Formic acid is used in leather tanning, textile dyeing and finishing.
2. Acetic acid is used in the manufacture of rayon and in plastics, in rubber and silk industries, in cooking and in vinegar (a 8-10% solution of acetic acid).
3. Benzoic acid and its salts are used as urinary antiseptics.
4. Formic acid can act as a reducing agent.


## Derivatives of Carboxylic acids

These are obtained when -OH group of carboxylic acids is replaced by Cl, NH<sub>2</sub>, OR' and OCOR and are called respectively acid chloride, acid amide, ester and acid anhydride.




## Properties of Acid Derivatives


### 1. Chemical reactions of acid halides



### 2. Chemical reactions of acid amides



### 3. Chemical reactions of ester



### 4. Chemical reactions of anhydrides

