Define environmental chemistry. Answer

Environmental chemistry is the study of chemical and biochemical processes occurring in nature. It deals with the study of origin, transport, reaction, effects, and fates of various chemical species in the environment.

Ouestion 14.2:

Ouestion 14.1:

Explain tropospheric pollution in 100 words.

Answer

Tropospheric pollution arises due to the presence of undesirable substances in the lowest

layer of the atmosphere. Oxides of sulphur, nitrogen, carbon, and hydrocarbons are the major gaseous pollutants. Oxides of sulphur (SO₂ and SO₃) and nitrogen (NO₂, NO) are produced as a result of \mathbb{C}

burning of fossil fuels (coal, automobile fuel). These oxides react with water in the

presence of atmospheric oxygen to form nitric acid (HNO₃) and sulphuric acid (H₂SO₄),

which leads to the formation of 'Acid rain'. $2SO_{2(g)} + O_{2(g)} + 2H_2O_{(l)} \longrightarrow 2H_2SO_{4(aq)}$ $4NO_{2(g)} + O_{2(g)} + 2H_2O_{(l)} \longrightarrow 4HNO_{3(aq)}$ Acid rain causes harm to agriculture, plants, and trees. It also leads to various respiratory ailments

respiratory ailments. Hydrocarbons are carbon and hydrogen containing compounds that burn to produce oxides of carbon. Hydrocarbons are carcinogenic and their products are also major pollutants. Carbon monoxide (CO) is poisonous in nature as it reacts with the

haemoglobin of blood, which can even result in death. Though carbon dioxide (CO_2) is not toxic in nature, yet it contributes towards global warming by trapping the reflected

IR rays. This results in the heating up of the Earth's atmosphere, thereby leading to the melting of icebergs and glaciers.

Particulates of smoke, dust, mist, and fume are harmful for human health as they are likely to block the nasal passage of a person, causing respiratory ailments. Smoke and fog combine to produce smog during a cool, humid day, thereby reducing visibility to a

large extent. Photochemical smog is formed due to the presence of PAN, ozone,

www.ncerthelp.com

Ouestion 14.3:

leads to the cracking of rubber and does damage to plants.

Carbon monoxide gas is more dangerous than carbon dioxide gas. Why?

Answer

Carbon dioxide (CO₂) and carbon monoxide (CO) gases are emitted during the

combustion of various fuels. Carbon monoxide is poisonous, whereas carbon-dioxide is non-toxic in nature.

Carbon monoxide is poisonous because it is capable of forming a complex with haemoglobin (carboxyhaemoglobin), which is more stable than the oxygen-haemoglobin complex. The concentration range of 3-4% of carboxyhaemoglobin decreases the

formaldehyde, and acrolein. It causes eve irritation, headaches, and chest pain. It also

oxygen-carrying capacity of blood. This results in headaches, weak eyesight, \geq nervousness, and cardiovascular disorders. A more increased concentration may even lead to death.

www.ncerthelp.com

Carbon dioxide is not poisonous. It proves harmful only at very high concentrations.

Question 14.4: List gases which are responsible for greenhouse effect.

Answer

The major greenhouse gases are: 1) Carbon dioxide (CO₂)

- 2) Methane (CH₄)
- 3) Water (H₂O)
- 4) Nitrous oxide (NO) 5) Ozone (O₃)
- 6) Chlorofluorocarbons (CFCs)

Question 14.5:

Answer

Statues and monuments in India are affected by acid rain. How?

Acid rain is a byproduct of various human activities that leads to the emission of oxides of sulphur and nitrogen in the atmosphere. These oxides undergo oxidation and then react with water vapour to form acids.

$$2SO_{2(g)} + O_{2(g)} + 2H_2O_{(l)} \longrightarrow 2H_2SO_{4(aq)}$$
$$4NO_{2(g)} + O_{2(g)} + 2H_2O_{(l)} \longrightarrow 4HNO_{3(aq)}$$

Acid rain causes damage to buildings and structures made of stone and metal. In India, limestone is a major stone used in the construction of various monuments and statues, including the Taj Mahal.

Acid rain reacts with limestone as:

$$CaCO_3 + H_2SO_4 \longrightarrow CaSO_4 + H_2O + CO_2$$

This results in the loss of lustre and colour of monuments, leading to their disfiguration.

Question 14.6:

What is smog? How is classical smog different from photochemical smog?

Answer

Smog is a kind of air pollution. It is the blend of smoke and fog. There are two kinds of smog:

a) Classical smog
b) Photochemical smog
The two smogs can be differentiated as follows: Classical smog Photochemical smog It occurs in a cool, humid It occurs in a dry, sunny climate. Occurrence climate. Smoke, PAN, acrolein, ozone, formaldehyde, fog, and Components sulphurdioxide. nitric oxide. Nature It is reducing in nature It is oxidizing in nature.

Ouestion 14.7:

Write down the reactions involved during the formation of photochemical smog.

Answer

Photochemical smog is formed as a result of the reaction of sunlight with hydrocarbons and nitrogen oxides. Ozone, nitric oxide, acrolein, formaldehyde, and peroxyacetyl nitrate (PAN) are common components of photochemical smog. The formation of

Burning of fossil fuels leads to the emission of hydrocarbons and nitrogen dioxide in the atmosphere. High concentrations of these pollutants in air results in their interaction with sunlight as follows:

$$NO_{2(g)} \xrightarrow{h\nu} NO_{(g)} + O_{(g)}$$

photochemical smog can be summarized as follows:

$$O_{(g)}$$
 + $O_{2(g)}$ \longleftrightarrow $O_{3(g)}$
 $O_{(g)}$ + $O_{(g)}$ \longleftrightarrow $O_{(g)}$ +

Nitrogen dioxide Nitric oxide $O_{(g)} + O_{2(g)} \longleftrightarrow O_{3(g)} + O_{2(g)} \longleftrightarrow O_{3(g)} + O_{2(g)} \longleftrightarrow O_{2(g)} + O_{2(g)}$ While ozone is toxic in nature, both NO₂ and O₃ are oxidizing agents. They react with the unburnt hydrocarbons in air to produce formaldehyde, PAN, and acrolein. $3CH_4 + 2O_3 \rightarrow 3CH_2 = O + 3H_2O$ Formaldehyde

Question 14.8:

What are the harmful effects of photochemical smog and how can they be controlled?

Answer

EFFECTS OF PHOTOCHEMICAL SMOG:

dryness, and various respiratory ailments.

Photochemical smog is oxidizing smog owing to the presence of NO₂ and O₃, causing corrosion of metals, stones, rubber, and painted surfaces. The other major components of photochemical smog are PAN, acrolein, and formaldehyde. Both PAN and ozone are eye irritants, while nitric oxide (formed from NO2) causes nose and throat irritation. At

higher concentrations, photochemical smog causes chest pain, headaches, throat

CONTROL MEASURES:

are decomposed by UV radiations. $CF_2Cl_{2(g)} \xrightarrow{UV} \dot{C}l_{(g)} + \dot{C}F_2Cl_{(g)}$

The chlorine free radical produced in reaction (iii) reacts with ozone as:

Photochemical smog results from the burring of fossil fuels and automobile fuels that emit NO₂ and hydrocarbons, which in turn form ozone, PAN, and other chemicals. The use of catalytic converters in automobiles is recommended to prevent the release of NO₂

Plantation of plants such as Pinus, Juniparur, Ouercus, Pyrus, and Vitis is also advised as

What are the reactions involved for ozone layer depletion in the stratosphere?

In the stratosphere, ozone is a product of the action of UV radiations on dioxygen as:

industries.

Once released CFCs mix with atmospheric gases and reach the stratosphere, where they

(iv) $\dot{C} l_{(g)} + O_{3(g)} \longrightarrow Cl \dot{O}_{(g)} + O_{2(g)}$

and hydrocarbons into the atmosphere.

Ouestion 14.9:

 $O_{2(g)} \xrightarrow{\text{UV}} O_{(g)} + O_{(g)}$

 $O_{2(g)} + O_{(g)} \leftarrow O_{3(g)}$

Answer

these plants have the capability to metabolize NO₂.

The $\operatorname{Cl}_{(g)}$ radicals further react with atomic oxygen to produce more chlorine radicals as: (v) $Cl\dot{O}_{(g)} + O_{(g)} \longrightarrow \dot{C}l_{(g)} + O_{2(g)}$

The regeneration of $^{\mathbf{Cl}_{(g)}}$ causes a continuous breakdown of ozone present in the stratosphere, damaging the ozone layer.

hypochlorous acid, which react further to give molecular chlorine. Molecular chlorine and

Ouestion 14.10:

What do you mean by ozone hole? What are its consequences?

Answer In Polar regions, stratospheric clouds provide the surface for chlorine nitrate and

HOCl are photolysed to give chlorine-free radicals.
$${\rm ClONO}_{2(g)} + {\rm H}_2{\rm O}_{(g)} {\longrightarrow} {\rm HOCl}_{(g)} + {\rm HNO}_{3(g)}$$

$$CIONO_{2(g)} + HCl_{(g)} \longrightarrow Cl_{2(g)} + HNO_{3(g)}$$

$$\begin{array}{c} \text{CIONO}_{2(g)} + \text{H2}_{2}\text{O}_{(g)} \longrightarrow \text{CI}_{2(g)} + \text{HNO}_{3(g)} \\ \text{HOCl}_{(g)} \stackrel{h\nu}{\longrightarrow} \dot{\text{O}}\text{H}_{(g)} + \dot{\text{CI}}_{(g)} \\ \text{CI}_{2(g)} \stackrel{h\nu}{\longrightarrow} \dot{\text{CI}}_{(g)} + \dot{\text{CI}}_{(g)} \\ \text{The chlorine-free radicals lead to the decomposition of ozone as:} \\ \dot{\dot{\text{CI}}_{(g)}} + O_{3(g)} \longrightarrow \text{CI}\dot{\text{O}}_{(g)} + O_{2(g)} \\ \text{Hence, a chain reaction is initiated. The chlorine-free radical is continuously regenerated, thereby depleting the ozone layer. This phenomenon is known as the as 'ozone hole'.} \end{array}$$

Effects of depletion of ozone layer The ozone layer protects the Earth from the harmful UV radiations of the sun. With the depletion of the layer, more radiation will enter the Earth's atmosphere. UV radiations

are harmful because they lead to the ageing of skin, cataract, skin cancer, and sunburns.

They cause death of many phytoplanktons, which leads to a decrease of fish productivity. Excess exposure may even cause mutation in plants.

Increase in UV radiations, decreases the moisture content of the soil and damages both plants and fibres.

Question 14.11:

What are the major causes of water pollution? Explain.

Answer www.ncerthelp.com

presence of several undesirable substances in water. Major water pollutants with their sources have been tabulated as follows:	
Pollutant	Source
Micro-organisms	Domestic sewage
Organic wastes	Domestic sewage, decaying animals and plants, animal excreta and waste, discharge from food processing industries

Water pollution arises as a result of several human activities, which leads to the

Plant nutrients Chemical fertilizers

heavy

Toxic

metals

Sediments

Pesticides

Strip mining and soil erosion

Chemical factories and industries

Radioactive Mining of uranium-containing minerals substances

Water used for cooling in industries

Heat

Roles played by major pollutants are: 1. Pathogens: These water pollutants include bacteria and other organisms. They enter

water from animal excreta and domestic sewage. Bacteria present in human excreta (for example, Escherichia coli and Streptococcus faecalis) cause gastrointestinal diseases.

run off. The presence of excess organic wastes in water decreases the amount of oxygen held by water. This decrease in the amount of dissolved oxygen inhibits aquatic life. 3. Chemical pollutants: These are water soluble chemicals like heavy metals such as

2. Organic wastes: These are biodegradable wastes that pollute water as a result of

Chemicals used for killing fungi, weed, insects

cadmium, mercury, nickel, etc. The presence of these chemicals (above the tolerance limit) can damage the kidneys, central nervous system, and liver.

Question 14.12:

Therefore, all industrial and chemical discharges should be made free from toxic metals before allowing them to enter a water body. The concentration of these pollutants should be checked regularly. Compost should be preferred over chemical fertilizers in gardens and agricultural fields to avoid harmful chemicals from entering ground water.

Answer Biochemical oxygen demand is the amount of oxygen required by bacteria to decompose

Ouestion 14.13:

suggest to control it?

contaminating the water and rendering it impure.

What do you mean by Biochemical Oxygen Demand (BOD)?

Answer

drinkina.

value of less than 5 ppm, whereas highly polluted water has a BOD value of 17 ppm or more.

organic matter in a certain volume of sample of water. Clean water would have a BODO

Have you ever observed any water pollution in your area? What measures would you

Water pollution arises as a result of various human activities. This includes discharges from wastewater treatment plants, run-off from agricultural fields, storm-water drainage, etc. Pollutants from these sources enter the water bodies, thereby

Industries and chemical factories discharge toxic, heavy metals such as Fe, Mn, Al, etc., along with organic wastes into water. Domestic sewage and animal excreta are also responsible for pathogenic contamination of water. These pollutants make water unfit for

Question 14.14:

Do you observe any soil pollution in your neighbourhood? What efforts will you make for controlling the soil pollution?

Answer

Major sources of soil pollution are industrial wastes and agricultural pollutants such as pesticides, fertilizers, etc.

It is very important to maintain the quality and fertility of soil to ensure and sustain the growth of plants and food crops.

www.ncerthelp.com

Pesticides are a mixture of two or more substances. They are used for killing pests. Pests include insects, plant pathogens, weeds, mollusks, etc., that destroy the plant crop and spread diseases. Aldrin and dieldrin are the names of some common pesticides Herbicides are pesticides specially meant for killing weeds. For example, sodium chlorate (NACIO₃), sodium arsenite (Na₃AsO₃), etc.

Question 14.16:

What do you mean by green chemistry? How will it help decrease environmentally pollution?

Answer

Ouestion 14.15:

What are pesticides and herbicides? Explain giving examples.

then, allowed to be dumped.

Answer

Answer

Green chemistry is a production process that aims at using the existing knowledge and

principles of chemistry for developing and implementing chemical products and processes to reduce the use and generation of substances hazardous to the environment The release of different harmful chemicals (particulates, gases, organic and inorganic

Insecticides like DDT are not soluble in water. For this reason, they remain in soil for a long time, contaminating the root crops. Pesticides like Aldrin and Dieldrin are nonbiodegradable and highly toxic in nature. They can enter the higher trophic levels through food chains, causing metabolic and physiological disorders. The same is true for

Hence, the best way to check soil pollution is to avoid direct addition of pollutants to the soil. Also, wastes should undergo proper treatment. They should be recycled and only

industrial wastes that comprises of several toxic metals like Pb, As, Hq, Cd, etc.

wastes) causes environmental pollution. In green chemistry, the reactants to be used in chemical reactions are chosen in such a way that the yield of the end products is up to 100%. This prevents or limits chemical pollutants from being introduced into the

environment. Through the efforts of green chemists, H₂O₂ has replaced tetrachlorethane

Question 14.17:

and chlorine gas in drying and bleaching of paper.

www.ncerthelp.com

atmosphere? Discuss.

Answer

Earth's most abundant greenhouse gases are CO₂, CH₄, O₃, CFCs, and water vapour.

These gases are present near the Earth's surface. They absorb solar energy that is radiated back from the surface of the Earth. The absorption of radiation results in the heating up of the atmosphere. Hence, greenhouse gases are essential for maintaining

In the absence of greenhouse gases, the average temperature of the Earth will decrease

drastically, making it uninhabitable. As a result, life on Earth would be impossible.

What would have happened if the greenhouse gases were totally missing in the earth's

of toxic dumping but you find an abundance of phytoplankton. Suggest a reason for the

Ouestion 14.18:

fish kill.

Answer

The amount of dissolved oxygen present in water is limited. The abundance of

phytoplanktons causes depletion of this dissolved oxygen. This is because, \square

As a result, the BOD level of water drops below 6 ppm, inhibiting the growth of fish and

A large number of fish are suddenly found floating dead on a lake. There is no evidence

phytoplanktons are degraded by bacteria present in water. For their decomposition, they require a large amount of oxygen. Hence, they consume the oxygen dissolved in water.

causing excessive fish-kill.

Question 14.19:

the temperature of the Earth for the sustenance of life.

How can domestic waste be used as manure?

Answer

Depending upon the nature of the waste, domestic waste can be segregated into two categories i.e., biodegradable and non-biodegradable. Biodegradable waste such as

leaves, rotten food, etc. should be deposited in land fills, where they get decomposed aerobically and anaerobically into manure. Non-biodegradable waste (which cannot be degraded) such as plastic, glass, metal scraps etc. should be sent for recycling.

Question 14.20:

Discuss the process in the light of bad odour, flies and recycling of wastes for a good produce.

Answer

For your agricultural field or garden you have developed a compost producing pit.

It is essential to take proper care of the compost producing pit in order to protect

ourselves from bad odour and flies.

It should be kept covered to minimize bad odour and prevent flies from entering it.

The recyclable waste should not be dumped in the compost producing pit. It should be sent to the industries through vendors for recycling.