Question 1:

In a cricket math, a batswoman hits a boundary 6 times out of 30 balls she plays.

Find the probability that she did not hit a boundary.

Answer:

Number of times the batswoman hits a boundary = 6

Total number of balls played = 30

 \therefore Number of times that the batswoman does not hit a boundary = 30 - 6 = 24

P (she does not hit a boundary) =	Number of times when she does not hit boundary
	Total number of balls played

 $=\frac{24}{30}=\frac{4}{5}$

Question 2:

1500 families with 2 children were selected randomly, and the following data were d:

re	CO	rc	le

,			
Number of families	475	814	211
Compute the probability of a fa	mily	chacar	at ra

Compute the probability of a family, chosen at random, having

(i) 2 girls (ii) 1 girl (iii) No girl

Number of girls in a family | 2

Also check whether the sum of these probabilities is 1.

Answer:

= 1500

Total number of families = 475 + 814 + 211

- (i) Number of families having 2 girls = 475

P₁ (a randomly chosen family has 2 girls) =
$$\frac{\text{Number of families having 2 girls}}{\text{Total number of families}}$$

= $\frac{475}{1500} = \frac{19}{60}$

(ii) Number of families having 1 girl = 814

 P_2 (a randomly chosen family has 1 girl) = $\frac{\text{Number of families having 1 girl}}{}$ Total number of families

(iii) Number of families having no girl = 211

Number of families having no girl P₃ (a randomly chosen family has no girl) = Total number of families


Sum of all these probabilities = $\frac{19}{100} + \frac{407}{100} + \frac{211}{100}$ $=\frac{475+814+211}{1500}$ $=\frac{1500}{1500}=1$

Therefore, the sum of all these probabilities is 1.

Question 3:

In a particular section of Class IX, 40 students were asked about the months of their

birth and the following graph was prepared for the data so obtained:



Find the probability that a student of the class was born in August.

Number of students born in the month of August = 6

Total number of students — 40

Total number of students = 40

 $P ext{ (Students born in the month of August)} = \frac{Number of students born in August}{Total number of students}$

Answer:

Three coins are tossed simultaneously 200 times with the following frequencies of

differe	different outcomes:					
	Outcome	3 heads	2 heads	1 head	No head	
	Frequency	23	72	77	28	

If the three coins are simultaneously tossed again, compute the probability of 2 heads coming up.

Answer:

Number of times 2 heads come up = 72

Total number of times the coins were tossed = 200

P(2 heads will come up) =	Number of times 2 heads come up
1 (2 heads will come up)	Total number of times the coins were tossed
	72 9
=	$\frac{1}{200} = \frac{1}{25}$

Question 5:

An organization selected 2400 families at random and surveyed them to determine a relationship between income level and the number of vehicles in a family. The information gathered is listed in the table below:

	Vehicles per family			
(in Rs)	0	1	2	Above 2

	2
	1
	25
	88
1	e probabi
ı	month and
	month and

(iii) earning less than Rs 7000 per month and does not own any vehicle.

535 + 29 + 1 + 2 + 469 + 59 + 25 + 1 + 579 + 82 + 88 = 2400

(iv) earning Rs 13000 – 16000 per month and owning more than 2 vehicles.

Number of total families surveyed = 10 + 160 + 25 + 0 + 0 + 305 + 27 + 2 + 1 + 100

(i) Number of families earning Rs 10000 - 13000 per month and owning exactly 2

25

27

29

59

0

160

305

535

469

16000 or more 579 82 1 Suppose a family is chosen, find th

10

0

1

2

Less than 7000

7000 - 10000

10000 - 13000

13000 - 16000

Answer:

(i) earning Rs 10000 - 13000 per (ii) earning Rs 16000 or more per month and owning exactly 1 vehicle.

vehicles = 29 $P = \frac{29}{2400}$ Hence, required probability,

Hence, required probability,

(v) owning not more than 1 vehicle.

(ii) Number of families earning Rs 16000 or more per month and owning exactly 1 vehicle = 579 $P = \frac{579}{2400}$

(iii) Number of families earning less than Rs 7000 per month and does not own any vehicle =
$$10$$

Hence, required probability,

2 vehicles = 25 $P = \frac{25}{2400} = \frac{1}{96}$ Hence, required probability

(iv) Number of families earning Rs 13000 - 16000 per month and owning more than

Hence, required probability,
$$2400-96$$

(v) Number of families owning not more than 1 vehicle = $10 + 160 + 0 + 305 + 1 + 100$

535 + 2 + 469 + 1 + 579 = 2062

Hence, required probability, $P = \frac{2062}{2400} = \frac{1031}{1200}$ Question 6:

A teacher wanted to analyse the performance of two sections of students in a

mathematics test of 100 marks. Looking at their performances, she found that a few students got under 20 marks and a few got 70 marks or above. So she decided to group them into intervals of varying sizes as follows: 0 - 20, 20 - 30... 60 - 70, 70 - 100. Then she formed the following table:

 Marks
 Number of student

Marks	Number of student
0 – 20	7
20 – 30	10
30 - 40	10
40 - 50	20
50 – 60	20
60 – 70	15
70 – above	8
Total	90
	0 - 20 20 - 30 30 - 40 40 - 50 50 - 60 60 - 70 70 - above

(i) Find the probability that a student obtained less than 20 % in the mathematics

- test.
- (ii) Find the probability that a student obtained marks 60 or above.

Answer:

Totalnumber of students = 90

 $P = \frac{7}{90}$

(i) Number of students getting less than 20 % marks in the test = 7

(ii) Number of students obtaining marks 60 or above = 15 + 8 = 23

 $P = \frac{23}{90}$

Question 7:

Hence, required probability,

Hence, required probability,

To know the opinion of the students about the subject statistics, a survey of 200 students was conducted. The data is recorded in the following table.

Opinion	Number of students
like	135
dislike	65
aisiike	00

Find the probability that a student chosen at random

(i) likes statistics, (ii) does not like it

Answer:

Total number of students = 135 + 65 = 200

- (i) Number of students liking statistics = 135
- P(students liking statistics) = $\frac{135}{200} = \frac{27}{40}$
- (ii) Number of students who do not like statistics = 65 P(students not liking statistics) = $\frac{65}{200} = \frac{13}{40}$

Question 8:

The distance (in km) of 40 engineers from their residence to their place of work were found as follows.

- 5 3 25 12 31 10 20 11
- 19 10 12 18 11 32 17

12 14 2 9 6 15 15 7 6 12

What is the empirical probability that an engineer lives:

- (i) less than 7 km from her place of work?
- (ii) more than or equal to 7 km from her place of work?

(iii) within
$$\frac{2}{2}$$
 km from her place of work?

Answer:

(i) Total number of engineers = 40

Number of engineers living less than 7 km from their place of work = 9

Hence, required probability that an engineer lives less than 7 km from her place of

$$P = \frac{1}{40}$$
 work, (ii) Number of engineers living more than or equal to 7 km from their place of work

= 40 - 9 = 31

Hence, required probability that an engineer lives more than or equal to 7 km from

her place of work,
$$\frac{1}{40}$$

(iii) Number of engineers living within
$$\frac{2}{2}$$
 km from her place of work = 0

Hence, required probability that an engineer lives within $^2\,\mathrm{km}$ from her place of work, P=0

Question 11:

Eleven bags of wheat flour, each marked 5 kg, actually contained the following weights of flour (in kg):

4.97 5.05 5.08 5.03 5.00 5.06 5.08 4.98 5.04 5.07 5.00

Find the probability that any of these bags chosen at random contains more than 5 kg of flour.

Answer:

Number of total bags = 11

Number of bags containing more than 5 kg of flour = 7

 $\label{eq:P} P = -\frac{1}{2}$ Hence, required probability,

Question 12:

Concentration of SO ₂ (in ppm)	Number of days (frequency)
0.00 - 0.04	4
0.04 - 0.08	9
0.08 - 0.12	9
0.12 - 0.16	2
0.16 - 0.20	4
0.20 - 0.24	2
Total	30

The above frequency distribution table represents the concentration of sulphur dioxide in the air in parts per million of a certain city for 30 days. Using this table, find the probability of the concentration of sulphur dioxide in the interval 0.12-0.16 on any of these days.

Answer:

Number days for which the concentration of sulphur dioxide was in the interval of 0.12-0.16=2

Total number of days = 30

	P	$r = \frac{2}{30}$	= -
Hence, required probability,	,	30	15

Question 13:

Blood group	Number of students
А	9
В	6
AB	3
0	12
Total	30

The above frequency distribution table represents the blood groups of 30 students of a class. Use this table to determine the probability that a student of this class, selected at random, has blood group AB.

Answer:

Number of students having blood group AB = 3

$$P = \frac{3}{20} = \frac{1}{10}$$

Total number of students = 30

Hence, required probability,
$$P = \frac{3}{30} = \frac{1}{10}$$